

An Overview on Staining Potential of Root Canal Irrigants and Medicaments in

¹Nitish Mathur, ²Kartik S Nath, ³Yusuf Md Nasir Khan, ⁴Anupam Singh, ⁵Tarun Nagpal, ⁶Prashant Kothari

Abstract--*Success of endodontic treatment depends upon eliminating microorganisms from the root canal. Thus, disinfection in endodontics is a critical step which relies on root canal irrigants and intracanal medicaments. This review aims to discuss the intrinsic and internalized staining potential of routinely used root canal irrigants and intracanal medicaments in endodontic practice. Keeping in mind the antimicrobial effect of irrigants and intracanal medicaments, the dentist should also consider their staining potential and measures to prevent the same. Thus, endodontic treatment should not focus only on biological and functional aspects, but also consider aesthetic consequences as well.*

Key words--*Disinfection, Intracanal Medicaments, Root canal irrigants, Staining potential*

I. INTRODUCTION

Tooth colour known to be a critical principle of smile aesthetics¹ has a major impact on person's psychosocial well-being and dental self-confidence.² It takes substantial amount of time and money to improve the appearance of a discoloured tooth.³ Since discolouration of tooth can be attributed to a variety of reasons, hence they are mainly categorized as extrinsic and intrinsic.^{4,5}

Extrinsic discolouration can occur either when staining compounds are absorbed in the pellicle layer or as a result of the interaction between them and tooth surface. Intrinsic discolouration can occur due to the structural changes within the bulk of tooth that could either be systemic or pulpal in origin.⁶

When the extrinsic stains are taken up by the tooth due to developmental defects, dental caries and restorations, it is termed as internalized tooth discoloration.⁷

The primary aim of endodontic treatment is to achieve maximum disinfection and to prevent reinfection of the root canal. Root canal irrigants and intracanal medicaments are the endodontic materials which serve this purpose by eliminating microbial load in the root canal.^{8,9} However, past studies indicate that endodontic materials can compromise aesthetics of tooth by inducing coronal discolouration^{10,11,12}

¹Department of Conservative Dentistry and Endodontics, Peoples Dental Academy, Bhopal, India. Email: nitishomr@gmail.com

²Assistant Professor, Department of Conservative Dentistry and Endodontics, Melaka Manipal Medical College, Manipal, affiliated to Manipal Academy of Higher Education, Karnataka, India. Email: kartik.nath@manipal.edu

³Consultant Orthodontist, Department of Orthodontics and Dentofacial Orthopaedics, Clove Dental, Bangalore, India. Email: yusufmknk@gmail.com

⁴Lecturer, Department of Oral and Maxillofacial Surgery, Manipal College of Dental Sciences, Manipal, affiliated to Manipal Academy of Higher Education, Karnataka, India. Email: anupam.singh@manipal.edu

⁵Consultant Oral and Maxillofacial Surgeon, Department of Oral and Maxillofacial Surgery, Link Hospital, Gwalior, India. Email: drtarunomfs@gmail.com

⁶Senior Lecturer, Department of Prosthodontics and Crown & Bridge, Chhattisgarh Dental College and Research Institute, Rajnandgaon, Chhattisgarh, India. Email: drprashantjainkothari@gmail.com

Besides having a proper knowledge of the antimicrobial effects of various root canal irrigants and intracanal medicaments, dentists should also be aware of their staining potential during usage in root canal therapy.

Thus, this review aims to describe the staining potential of routine root canal irrigants and intracanal medicaments used in endodontics

II. ROOT CANAL IRRIGANTS

Sodium hypochlorite (NaOCl)

Sodium hypochlorite (NaOCl) is the most commonly recommended endodontic irrigant as it possess high tissue dissolving property and wide-spectrum antimicrobial activity.^{13,14} Saponification reaction, amino acid neutralization and chloramination reactions are responsible for its pulp dissolution.¹⁵ Moreover, sodium hypochlorite is a bleaching agent too and will not be causing discoloration on its own. However, on its contact with the red blood corpuscles and its high tendency to crystallize on the dentinal walls may produce discoloration which will be difficult to remove it from the canal.¹⁶

NaOCl and Chlorhexidine (CHX) solution

An irrigation regimen combining NaOCl with CHX solution is used to enhance the antimicrobial properties.¹⁷ This combination however forms a dark brown precipitate which can compromise the aesthetics by staining dentin.¹⁸ The precipitate acts a residual film and may get adhered on the dentinal walls, floor of the pulp chamber and apart from staining, it may also affect the diffusion of root canal irrigants and medicaments into the dentin. Moreover, it will also interfere with the root canal obturation leading to a failure.^{19,20}

Examination of the dark brown precipitate using X-ray photoelectron spectroscopy (XPS) and time-of-flight secondary ion mass spectrometry (TOF-SIMS) reveals that it contains a significant amount of parachloroaniline (PCA) which is carcinogenic in nature.²¹

The precipitate formation can be prevented by including a thorough intermediate flush in the irrigation protocol with the solutions such as saline or sterile distilled water, followed by drying of the canal before the next solution is used.²²

NaOCl and MTAD

MTAD-Mixture of tetracycline isomer, citric acid and detergent has been shown to overcome the lack of antimicrobial substantivity and inability to remove smear layer of NaOCl in clinical settings.²³ For an optimum smear layer removal along with superior antimicrobial sensitivity, an irrigation protocol with 1.3% NaOCl for 20 minutes, followed by the irrigation of MTAD for 5 minutes has been proposed.²⁴ The high affinity of doxycycline to the dentin makes it a potential staining agent present in MTAD.^{25,26}

But since MTAD is delivered for a very short duration in the root canal and there is no usual exposure of canals to light, therefore the adult onset tetracycline tooth staining is not seen.²⁷

However, a chemical reaction between NaOCl and MTAD leads to the production of brown solution in root canals.²⁸ This reaction might be caused by the dentinal absorption and release of doxycycline present in MTAD with the exposure to final rinse of NaOCl.²⁸

A study also reports the formation of yellow precipitate on the walls of root canal when MTAD is used as a final rinse. In the same study, when MTAD is used as a final rinse following the initial rinse of 1.3% NaOCl, the light exposed root dentinal walls were red-purple stained due to photooxidation by NaOCl.²⁹ To prevent the oxidation effect of NaOCl and photodegradation of the doxycycline that is present in MTAD, an intermediate flush with sodium ascorbate is recommended which acts as a reducing agent.³⁰

CHX and EDTA

In routine endodontic therapy, 17% EDTA is used as a chelating agent which serves the purpose of smear layer removal and making the dentin soft for better instrumentation.^{31,32}

A white to pink precipitate has been reported to form on its interaction with CHX.^{33,34} The precipitate is formed as a result of electrostatic neutralization of the cationic chlorhexidine by the anionic EDTA.³⁴ The resultant precipitate doesn't show any significant amounts of p-chloroaniline unlike the interaction between NaOCl and CHX.³⁴

To prevent the outcome of above-mentioned combination of irrigants, Maleic acid can be substituted for EDTA, as it has been proven a better agent in smear layer removal which does not cause any precipitation and discoloration on its association with CHX.³⁵

Silver diamine fluoride

3.8% Silver diamine fluoride has shown to be used as a potential antimicrobial root canal irrigant against *E. faecalis* biofilm. In the same study, silver salts were seen occluding the dentinal tubules.³⁶ The formation of silver phosphate layer and silver sulphide precipitation on dentin is attributed to the black staining potential of silver diamine fluoride.^{37,38}

Herbal irrigants

Green tea extract

Green tea polyphenols, prepared from the young shoots of the tea plant *Camellia sinensis* has shown statistically significant antibacterial activity against *E. faecalis* biofilm formed on tooth substrate.³⁹ However, it should be noted that the tannins present in the green tea can cause staining of teeth.⁴⁰

Triphala juice

Triphala comprises dried and powdered fruits derived from three medicinal plants *Terminalia bellerica*, *Terminalia chebula*, and *Embllica officinalis*.⁴¹ This formulation is effective against *Faecalis* and the citric acid in triphala also removes the smear layer, making it a potential root canal irrigant.³⁹ The dentinal discoloration is observed following the use of commercially prepared triphala juice which can be attributed to its natural pigment or due to the presence of additives.⁴²

III. INTRACANAL MEDICAMENTS

Calcium Hydroxide based medicaments

Calcium hydroxide has noteworthy properties such as antimicrobial activity, hard tissue barrier formation, arrest of resorption all of which makes it to be used widely as an effective intracanal medicament and as a material of choice in various clinical scenarios.^{43,44} Some studies have reported tooth staining, although insignificant, following its application as an intracanal medicament.^{43,45,46,47} This staining can be attributed to the radiopacifiers present in the calcium hydroxide based intracanal medicaments.^{45,48}

Tetracycline based medicaments

Triple antibiotic paste (containing ciprofloxacin, metronidazole, and minocycline) is the most common intra-canal medicament that can cause tooth staining if they are not completely removed from the access cavity at a level coronal to the gingival margin.⁴⁹ Tooth staining has been reported in tetracycline based endodontic materials in the effect of light through a photoinitiated reaction as stated before.⁴⁹ However a significant staining due to minocycline is also seen in the absence of light which is due to its bond with the calcium of dentin forming insoluble complex.⁴⁹

Ledermix paste contains a combination of a tetracycline antibiotic agent, 3.2% demeclocycline hydrochloride and a steroid agent, 1% triamcinolone acetonide in a polyethylene glycol base.⁵⁰ Effect of light, oxidation reaction and presence of moisture are the key mechanisms which are attributed to the staining potential of demeclocycline present in the paste.⁵⁰

To prevent the discoloration, doxycycline hylate must be substituted to demeclocycline hydrochloride, keeping the other component of paste the same.⁵¹ Incorporation of ascorbic acid, an oxidising agent into the paste or rinsing with it before placement of the medicament can also be an approach to prevent discoloration.⁵² Moreover, prevention of moisture and oxygen is the most important factor in preventing the tooth staining.

Iodoform based medicaments

A study evaluated the clinical success of pulpally treated primary incisors revealed a yellowish discoloration which was clearly seen through the coronal composite restoration. The iodoform paste used as an intracanal medicament in the study was concluded to be the reason for this yellow staining.⁵³

Formocresol based medicaments

The small molecular composition of formocresol aids its penetration into wide dentinal tubules, seen especially in young patients which reports to cause coronal discoloration.^{55,54} Gingival necrosis and crestal bone sequestration are also seen following the use of formocresol based medicaments.⁵⁵

IV. CONCLUSION

Almost all the irrigants and intracanal medicaments used in modern endodontics have shown to cause tooth staining. Therefore, keeping in mind, the staining potential of these agents during endodontic therapy, the dentist

must know the preventive measures and treatment options for the same. The endodontic therapy should not only focus on the biological and functional aspects, but should also consider the aesthetic outcome of procedure.

REFERENCES

1. Machado Andre Wilson. 10 commandments of smile esthetics. Dental Press J. Orthod. [Internet]. 2014 Aug [cited 2020 Mar 30]; 19(4): 136-157.
2. Afroz, S., Rathi, S., Rajput, G. *et al.* Dental Esthetics and Its Impact on Psycho-Social Well-Being and Dental Self Confidence: A Campus Based Survey of North Indian University Students. *J Indian Prosthodont Soc* 13, 455–460 (2013).
3. Sulieman MA (2008) An overview of tooth-bleaching techniques: chemistry, safety and efficacy. *Periodontology* 2000 48, 148–69.
4. Hattab FN, Qudeimat MA, al-Rimawi HS (1999) Dental discoloration: an overview. *Journal of Esthetic Dentistry* 11, 291–310
5. Plotino G, Buono L, Grande NM, Pameijer CH, Somma F (2008) Nonvital tooth bleaching: a review of the literature and clinical procedures. *Journal of Endodontics* 34, 394–407
6. Watts A, Addy M (2001) Tooth discolouration and staining: a review of the literature. *British Dental Journal* 190, 309–16.
7. Sulieman M (2005) An overview of tooth discoloration: extrinsic, intrinsic and internalized stains. *Dental Update* 32, 463–4. 466-8, 471.
8. Kumar A, Tamanna S, Iftekhar H. Intracanal medicaments – Their use in modern endodontics: A narrative review. *J Oral Res Rev* 2019;11:94-9
9. Kandaswamy D, Venkateshbabu N. Root canal irrigants. *J Conserv Dent.* 2010 Oct;13(4):256-64. doi: 10.4103/0972-0707.73378. PMID: 21217955; PMCID: PMC3010032.
10. Krastl G, Allgayer N, Lenherr P, Filippi A, Taneja P, Weiger R. Tooth discoloration induced by endodontic materials: a literature review. *Dental traumatology.* 2013 Feb;29(1):2-7.
11. Chahande RK, Patil SS, Gade V, Meshram R, Chandhok DJ, Thakur DA. Spectrophotometric analysis of crown discoloration induced by two different sealers: An *In vitro* study. *Indian J Dent Res* 2017;28:71-5
12. Gutierrez JH, Guzmán M (1968) Tooth discoloration in endodontic procedures. *Oral Surgery Oral Medicine Oral Pathology* 26, 706–11.
13. Naenni N, Thoma K, Zehnder M. Soft tissue dissolution capacity of currently used and potential endodontic irrigants. *J Endod* 2004;30:785-787.
14. Zehnder M, Kosicki D, Luder H, Sener B, Waltimo T. Tissue-dissolving capacity and antibacterial effect of buffered and unbuffered hypochlorite solutions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2002;94:756-762
15. Estrela C, Estrela CRA, Barbin EL, Spanó JCE, Marchesan MA, Pécora JD. Mechanism of action of sodium hypochlorite. *Braz Dent J.* 2002; 13(2):113-7.
16. Gutierrez JH, Guzmán M (1968) Tooth discoloration in endodontic procedures. *Oral Surgery Oral Medicine Oral Pathology* 26, 706–11.
17. Kuruvilla JP, Kamath P. Antimicrobial activity of 2.5% sodium hypochlorite and 0.2% chlorhexidine gluconate separately and combined, endodontic irrigants. *J Endod* 1998;24:472–6.
18. Marchesan MA, Pasternak JRB, Afonso MMF, et al. Chemical analysis of the flocculate formed by the association of sodium hypochlorite and chlorhexidine. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;103:e103–5.
19. Vivacqua-Gomes N, Ferraz CC, Gomes BP, Zaia AA, Teixeira FB, Souza-Filho FJ (2002) Influence of irrigants on the coronal microleakage of laterally condensed gutta-percha root fillings. *International Endodontic Journal* 35, 791–5.
20. Akisue E, Tomita VS, Gavini G, Poli de Figueiredo JA (2010) Effect of the combination of sodium hypochlorite and chlorhexidine on dentinal permeability and scanning electron microscopy precipitate observation. *Journal of Endodontics* 36, 847–50.
21. Basrani BR, Manek S, Sodhi RN, Fillery E, Manzur A (2007) Interaction between sodium hypochlorite and chlorhexidine gluconate. *Journal of Endodontics* 33, 966–9.
22. Krishnamurthy S, Sudhakaran S (2010) Evaluation and prevention of the precipitate formed on interaction between sodium hypochlorite and chlorhexidine. *Journal of Endodontics* 36, 1154–7.
23. Torabinejad M, Khademi AA, Babagoli J, et al. A new solution for the removal of the smear layer. *J Endod* 2003;29:170–5.
24. Torabinejad M, Cho Y, Khademi AA, Bakland LK, Shabahang S. The effect of various concentrations of sodium hypochlorite on the ability of MTAD to remove the smear layer. *J Endod* 2003;29:233–9.

25. Weyman J. The clinical appearances of tetracycline staining of the teeth. *Br Dent J* 1965;289-91.
26. Beltz RE, Torabinejad M, Poursmail M. Quantitative analysis of the solubilizing action of MTAD, sodium hypochlorite, and EDTA on bovine pulp and dentin. *J Endod* 2003;29:334–7
27. Park DS, Torabinejad M, Shabahang S. The effect of MTAD on the coronal leakage of obturated root canals. *J Endod* 2004;30:890–2.
28. Torabinejad M, Cho Y, Khademi AA, Bakland LK, Shabahang S. The effect of various concentrations of sodium hypochlorite on the ability of MTAD to remove the smear layer. *J Endod* 2003;29:233–9.
29. Tay FR, Mazzoni A, Pashley DH, Day TE, Ngoh EC, Breschi L (2006a) Potential iatrogenic tetracycline staining of endodontically treated teeth via NaOCl/MTAD irrigation: a preliminary report. *Journal of Endodontics* 32, 354–8
30. Nassar M, Awawdeh L, Jamleh A, Sadr A, Tagami J (2011) Adhesion of Epiphany self-etch sealer to dentin treated with intracanal irrigating solutions. *Journal of Endodontics* 37, 228–30
31. Sayin TC, Serper A, Cehreli ZC, Otlu HG. The effect of EDTA, EGTA, EDTAC, and tetracycline-HCl with and without subsequent NaOCl treatment on the microhardness of root canal dentin. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;104:418–24
32. Khedmat S, Shokouhinejad N. Comparison of the efficacy of three chelating agents in smear layer removal. *J Endod* 2008;34:599–602
33. Gonzalez-Lo'pez S, Camejo-Aguilar D, Sanchez-Sanchez P, Bolanos-Carmona V (2006) Effect of CHX on the decalcifying effect of 10% citric acid, 20% citric acid, or 17% EDTA. *Journal of Endodontics* 32, 781–4
34. Rasimick BJ, Nekich M, Hladek MM, Musikant BL, Deutsch AS (2008) Interaction between chlorhexidine digluconate and EDTA. *Journal of Endodontics* 34, 1521–3.
35. Ballal NV, Moorkoth S, Mala K, Bhat KS, Hussien SS, Pathak S (2011) Evaluation of chemical interactions of maleic acid with sodium hypochlorite and chlorhexidine gluconate. *Journal of Endodontics* 37, 1402–5.
36. Hiraishi N, Yiu CK, King NM, Tagami J, Tay FR. Antimicrobial Efficacy of 3.8% silver diamine fluoride and its effect on root dentin. *J Endod.* 2010;36:1026–9.
37. Patel J, Anthonappa RP, King NM. Evaluation of the staining potential of silver diamine fluoride: in vitro. *International journal of paediatric dentistry.* 2018 Sep;28(5):514-22.
38. Duangthip D, Wong MCM, Chu CH, Lo ECM. Caries arrest by topical fluorides in preschool children: 30-month results. *J Dent* 2018; 70: 74–79.
39. Prabhakar J, Senthilkumar M, Priya MS, Mahalakshmi K, Sehgal PK, Sukumaran VG. Evaluation of antimicrobial efficacy of herbal alternatives (*Triphala* and Green Tea Polyphenols), MTAD, and 5% sodium hypochlorite against *Enterococcus faecalis* Biofilm Formed on Tooth Substrate: An *In vitro* Study. *J Endod.* 2010;36:83–6
40. George J, Mathews T A. Spectrophotometric Analysis of Tooth Discoloration after Bracket Debonding Induced by Coffee and Tea Grown in South India: An *In vitro* Study. *J Interdiscip Dentistry* 2018;8:1-4
41. Jagetia GC, Baliga MS, Malagi KJ, Kamath SM. The evaluation of the radioprotective effect of Triphala (an Ayurvedic rejuvenating drug) in the mice exposed to radiation. *Phytomedicine.* 2002;9:99–108.
42. Choudhary E, Indushekar KR, Saraf BG, Sheoran N, Sardana D, Shekhar A. Exploring the role of *Morinda citrifolia* and Triphala juice in root canal irrigation: An ex vivo study. *Journal of conservative dentistry: JCD.* 2018 Jul;21(4):443.
43. Thomson AD, Athanassiadis B, Kahler B, Walsh L. Tooth discoloration: staining effects of various sealers and medicaments. *Aust Endod J.* 2012;38:2–9.
44. Kim ST, Abbott PV, McGinley P. The effects of Ledermix paste on discoloration of immature teeth. *Int Endod J.* 2000;33:233–237.
45. Lenherr P, Allgayer N, Weiger R, Filippi A, Attin T, Krastl G. Tooth discoloration induced by endodontic materials: a laboratory study. *Int Endod J.* 2012;45:942–949.
46. Ordinola-Zapata R, Bramante CM, Minotti PG, Cavenago BC, Garcia RB, Bernardineli N, Jaramillo DE, Hungaro Duarte MA. Antimicrobial activity of triantibiotic paste, 2% chlorhexidine gel, and calcium hydroxide on an intraoral-infected dentin biofilm model. *J Endod.* 2013;39:115–118.
47. Rahimi S, Janani M, Lotfi M, Shahi S, Aghbali A, Vahid Pakdel M, Salem Milani A, Ghasemi N. A review of antibacterial agents in endodontic treatment. *Iran Endod J.* 2014;9:161–168.
48. Almyroudi A, Mackenzie D, McHugh S, Saunders WP. The effectiveness of various disinfectants used as endodontic intracanal medications: an *in vitro* study. *J Endod.* 2002;28:163–167.
49. Kim JH, Kim Y, Shin SJ, Park JW, Jung IY (2010a) Tooth discoloration of immature permanent incisor associated with triple antibiotic therapy: a case report. *Journal of Endodontics* 36, 1086–91
50. Athanassiadis B, Abbott P.V., Walsh LJ, The Use Of Calcium Hydroxide, Antibiotics And Biocides As Antimicrobial Medicaments In Endodontics, *Australian Dental Journal,* (52), 2007, 64-82.

51. Chen BK, George R, Walsh LJ. Root discolouration following short-term application of steroid medicaments containing clindamycin, doxycycline or demeclocycline. *Aust Endod J.* 2012;38(3):124–8.
52. Bowles WH. Protection against minocycline pigment formation by ascorbic acid (vitamin C) *J Esthet Dent.* 1998;10(4):182–6.
53. Ari Kupietzky, William F.Waggoner, Jon Galea. The clinical and radiographic success of bonded resin composite strip crowns for primary incisors. *Pediatric dentistry.* 25: 6.2003. 577-581
54. Dankert J, Gravenmade EJ, Wemes JC (1976) Diffusion of formocresol and glutaraldehyde through dentin and cementum. *Journal of Endodontics* 2, 42–6
55. Cambrozzi JV, Greenfeld RS (1983) Necrosis of crestal bone related to the use of excessive formocresol medication during endodontic treatment. *Journal of Endodontics.*