TOOTH REGENERATION - A REVIEW

Ashwini.K¹, Dr.Jayalakshmi Somasundaram², Vinay Sivaswamy³

Abstract

Introduction: Dental problems caused by caries, PDL disease, and injury to the tooth compromise oral health issues resulting in loss of tooth thereby affecting the quality of human life. There are several artificial therapies introduced into dentistry to compensate for the loss of tooth structure and avulsed tooth structures ie) inorganic fillings, RCT, RPD, FPD, CD, and even implants. They tend to fail at times. The current advances in dentistry include regenerative therapy, stem cell biotechnology, and tissue engineering for 3D scaffolds. In this review, we focus on the recent findings and technologies relevant to whole teeth regenerative therapy.

Materials and methods: The articles and journals pertaining to this topic were searched over Pubmed, Google Scholar, Semantic Scholar, MeSH core, Elsevier, and Medline. Articles related to tooth regeneration, stem cell-based regeneration, and scaffolds were included. Wherein articles irrelevant to these topics are excluded. The data was collected from 2000 to 2020.

Discussion: Regenerative therapies for the whole tooth regeneration is a novel therapeutic concept. Dental stem cells and activating cytokines have a candidate approach for tooth regeneration and potential to differentiate tooth regeneration in vitro or in vivo and forms a functional tooth.3D tissue engineering for the bioengineered scaffolds regenerates organogenesis. They have several biomedical applications and have high bioactivity. A bioengineered tooth was also able to perform the same functions as a physiological tooth. The stem cells play an efficient role in the formation of tooth structures too.

Conclusion: Though tooth embryogenesis takes a longer duration, it is highly effective and promotes mental well being. There is also a huge potential for 3D printing of tissue engineering. Tooth regenerative therapy is actually a future regenerative technology.

Keywords: whole tooth regeneration, stem cells, 3D scaffold bioengineering

Introduction

The tooth is a very complex structure which has a self-limiting and healing capacity(1). It is an ectodermal organ and is composed of both hard and soft tissues-enamel, dentin, cementum, pulp, and periodontium(2). Tooth loss is commonly seen due to oral diseases like PDL disease, caries, traumatic injuries, etc. (3) and pathological causes which lead to physical and mental suffering that lower an individual's quality of life (4). There are several artificial therapies that compensate for the loss of the tooth structure and avulsed tooth structures(5) ie) RCT, FPD, RPD, CD, inorganic restorations and fillings, implants, etc(6)(7)(8)(9).

¹ Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77, Tamil Nadu, India, Email: <u>151801032@gmail.com</u>

² Corresponding author: Chief Scientist, White Lab-Material research center, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77, Tamil Nadu, India, Email: jayalakshmisomasundaram@saveetha.com

³ Senior Lecturer, Department of Prosthodontics and Implantology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77, Tamil Nadu, India, Email: <u>vinay.sdc@saveetha.com</u>

They provide and compensate for the loss but it is not said to be effective all the time (10,11). They tend to fail over a period of time. Further technological developments have been introduced to biologically repair and regenerate the tooth which is damaged (12). They are proven to be a novel therapeutic method for future dentistry (13).

One among them is the stem cell-based tissue engineering(14). They are isolated from the somatic stem cells in human dental tissues and children's embryonic cells. Their common lineage is that they are derived from the neural crest cells. The mesenchymal stem cells have several properties like marker genes. They also have cell-activating cytokines that differentiate tooth tissues in vitro and in vivo. Stem cells have the property and capacity to regenerate missing tissues or structures. They are undifferentiated.

The other therapeutic technology is the bioengineered 3D scaffolds(15). This tissue engineering is an emerging field of medicine that requires the combination of stem cells, biodegradable materials, and signaling molecules. The contribution of scaffolds is indispensable as they serve as carriers to facilitate the delivery of stem cells or growth factors at the receptor site. Scaffolds regenerate organogenesis and have several biomedical applications. They exhibit high bioactivity and have specific porosity.

A lot of focus has been made only in caries management(16)(17)(18)(19)(20)(19,21). There is only limited research focusing on tooth regeneration which is the future of modern dentistry. In this review, we mainly focus on the recent findings and technologies under tooth regenerative therapy.

MATERIALS AND METHODS:

The articles and journals pertaining to this topic were searched over Pubmed, Google Scholar, Semantic Scholar, MeSH core, Elsevier, and Medline. Articles related to tooth regeneration, stem cell-based regeneration, and scaffolds were included. Wherein articles irrelevant to these topics are excluded. The data was collected from 2000 to 2020.

DISCUSSION:

Regrowing a tooth:

A tooth is a complex organ consisting of soft connective tissue and mineralized hard tissues(1). Enamel is the highest mineralized tissue and is characterized by the absence of the cells. It is the first hardest barrier against the environment. The ameloblasts during enamel formation undergo programmed cell death during maturation. Thus once damaged enamel is prone to caries, trauma, etc. Artificial therapies like restoration provide only short term relief. Further developments in dentistry include tooth regenerative therapies with a whole tooth replacement which is one of the most emerging techniques in the world. Replacing and repairing enhances tissue or organ functions(22).

Stem cell-based tooth regeneration:

Stem cells are undifferentiated cells, capable of renewing themselves with the capacity to produce different cell types to regenerate missing tissues or structures(23)(24). Their microenvironment, composed of heterologous cell types, extracellular matrix, and soluble factors, enables them to maintain their stemness(25). Because of their unique properties, stem cells have the potential to be important in tissue engineering strategies for the regeneration of diseased, damaged, and missing tissues and even organs. In general, stem cells can be divided into three main types: ESCs that are derived from embryos; adult stem cells that are derived from adult tissue; and iPS cells that are generated artificially by reprogramming adult somatic cells so that they behave like ESCs. In this section, we outline recent results obtained using ESCs and adult stem cells for tooth regeneration. But the common lineage between both is that they are derived from the neural crest cells. The generic mesenchymal stem cells have properties like marker genes(26). They possess activating cytokines to differentiate into tooth tissues in both in vitro and in vivo conditions(14)(27).

International Journal of Psychosocial Rehabilitation, Vol. 24, Issue 02, 2020 ISSN: 1475-7192

Embryonic stem cells:

ESCs are pluripotent stem cells derived from the undifferentiated inner cell mass of the blastocyst. They continue to grow indefinitely in an undifferentiated diploid state and cultured in optimal conditions in the presence of a feeder layer and leukemia inhibitory factor. They can be differentiated into derivatives of all three primary germ layers: ectoderm, endoderm, and mesoderm. Because of the pluripotency of ESCs, it has a wide future clinical application(28)(29).In dentistry, ESCs have been used for oral tooth regeneration, including mucosa, alveolar bone, and periodontal tissue regeneration.ESCs expressed the unique set of genes for odontogenic mesenchymal cells, such as Lhx7, Msx1, and Pax9, suggesting that ESCs can respond to inductive signals from the embryonic dental epithelium. Although these approaches have the potential to be useful for tooth regeneration and for understanding basic tooth development, it will be necessary to address several major issues before they can be implemented in clinical practice when transplanted like the ethical issues regarding the use of embryos and allogeneic immune rejection.

Adult stem cells:

Adult stem cells have been identified in many tissues and organs and have been shown to undergo self-renewal, to differentiate for the maintenance of normal tissue, and to repair injured tissues(30).

Dental pulp stem cells:

When there is severe tooth damage that penetrates both enamel and dentin, the pulp produces a natural repair process and produces odontoblast. This proves there is a high possibility that tooth pulp has stem cells. The DPSCs (Dental pulp stem cells) are isolated from the human pulp(31)(32). They were first isolated from the 3rd molars and exhibit high proliferation and produce densely calcified nodules(33). They have the ability to regenerate neural tissues and dental pulp(34). They also differentiate into other mesenchymal cell derivatives in vitro. They have also demonstrated de novo regeneration of pulp in the emptied root canal. Clinical trials were carried in mice where DPSC and SCAP were seeded into D-L, lactide scaffold, and inserted into canal space. It is then transplanted subcutaneously to mice. The analysis after 3-4 months showed a pulp-like tissue with well-established vascularisation, Continuous layer of mineralized tissue was also seen on the walls of the canal(35). It is proposed that dental pulp stem cells (DPSCs) can develop Induced Pluripotent Stem Cells (iPSCs) which can be used for therapies of various diseases(36).

Stem cells from human exfoliated deciduous teeth:

MSC-like cells have also been isolated from the dental pulp of human deciduous teeth [stem cells from human exfoliated deciduous teeth (SHEDs)].SHEDs have the ability to differentiate in vitro to neuron-like cells, odontoblasts, osteoblasts, and adipocytes show higher proliferation rates and increased numbers of population doublings compared with DPSCs and can form spherical aggregations. When these cells are transplanted mixed with HA/TCP in vivo, they can form dentin and bone but not dentin–pulp complexes. A comparison of the gene expression profiles of DPSCs and SHEDs demonstrated that 4386 genes were differentially expressed by two-fold or more(37).In addition to genes, FGF, transforming growth factor (TGF)- β , and collagen I and III showed a higher level of gene expression in SHEDs than in DPSCs(38). SHEDs could be the ideal source of stem cells for repairing damaged teeth or for the induction of bone formation.

Periodontal stem cells:

PDL is a fibrous connective tissue containing specialized cells(39). They were recognized to contain a population of progenitor cells identified as periodontal ligament stem cells(40). They are capable of differentiating along mesenchymal cell lineages. It maintains PDL cell numbers.

Stem cells from the apical papilla:

Stem cells from the apical papilla (SCAPs) are found in the papilla tissue in the apical part of the roots of developing teeth.SCAPs are found in the third molars and teeth with open apices. These cells have the potential to differentiate into osteoblasts, odontoblasts, and adipocytes and show higher rates of proliferation in vitro

compared with DPSCs(41). Transplantation of SCAPs and periodontal ligament stem cells (PDLSCs) into tooth sockets of minipigs allowed the formation of dentin and periodontal ligament(42).

Dental follicle stem cells:

DFSCs have also been isolated from the follicles of developing third molars(43,44). They can differentiate into osteoblasts, adipocytes, and nerve-like cells in vitro and form cementum and periodontal ligament in vivo and form cementum and periodontal ligament in vivo(43)(45).

Future therapeutic approaches for the restoration of damaged dentin, pulp, cementum, and periodontal ligaments may make use of autologous stem cells such as DPSCs, SHEDs, SCAPs, and DFSCs that have been stored after removal from the patient.

ERMs and HERS:

The epithelial rests of Malassez (ERMs) are quiescent epithelial remnants of Hertwig's root sheath (HERS) that remain in the adult tooth and play a role in cementum repair and regeneration(46). A recent study demonstrated that ERMs contains a unique population of stem cells that are capable of undergoing epithelial-mesenchymal transition and differentiate into diverse lineages indicative of mesodermal and ectodermal origin, including bone, fat, and cartilage as well as neuron-like cells. In addition, ERMs can be induced to form enamel-like tissues after transplantation into an athymic rat omentum with primary dental pulp cells, suggesting that the stem cells in ERMs may be able to regenerate enamel(47).

Whole tooth regeneration:

Regenerating a whole tooth is the main objective of regenerative medicine(48). It increases the life expectancy in patients(49). It replaces the current state of the art replacement therapy namely dental implant. The clinical approach includes somatic stem cell usage. The formation of tooth germ is combined with inductive embryonic epithelial oral cells(15). The postnatal pulp cells are not suitable for tooth engineering. The induced pluripotent stem cells embryonic stem cells are the alternatives of tooth germ(49),(50). PLSC's induce the production of iPSCs(48,51). The decellularized tooth buds differentiate iPSCs to form tooth germ cells and promote tooth regeneration therapies.

3D printing technology:

Tissue engineering is an emerging field of medicine that is based on combining stem cells, growth factors, and scaffolds for regenerating lost tissues(15). The contribution of scaffolds is incredible as it serves as a carrier for the delivery of stem cells or growth factors at the local receptor site. Scaffold biomaterials can be natural or synthetic, rigid or non-rigid and degradable or non-degradable. Scaffolds are highly bioactive and have specific porosity(52). It has both soft and hard tissue components(53). They promote bone, PDL, and cementum formation. A synthetic scaffold is most accepted and it is predesigned(54). They act as a role guiding and assisting ECM(55). The clinical experiments include using dental epithelial and pulpal mesenchymal tissues which are seeded in the mixture of heterogeneous single cells into a tooth-shaped biodegradable polymer scaffold. These cell scaffolds were implanted into the host's body to receive sufficient oxygen and nutrients. After 25-30 weeks, the implanted scaffolds resembled like the crowns of natural teeth. It takes longer duration and the correct arrangement as of a natural tooth(56).

Future scope:

The stem cell-based regeneration provides a good quality of repair and has high efficiency. Bioengineering regulates tissue engineering scaffolds and cytokines morphogenesis(15). They have high biocompatibility and mechanical properties.

International Journal of Psychosocial Rehabilitation, Vol. 24, Issue 02, 2020 ISSN: 1475-7192

CONCLUSION:

A bioengineered tooth is also found to perform the same functions as a physiological tooth ie) mastication. It provides aesthetics and promotes confidence and mental well being. There is a huge potential in 3D printing because of its advantages like flexibility and internal porous structure(57). In the field of stem cell tooth regeneration, it provides an attractive alternative to traditional and current practices for the replacement of missing teeth, such as implants and classic procedures based on synthetic materials. Because of rapidly increasing research efforts and progress, it is anticipated that clinically satisfactory functional tooth regeneration will be available in the near future(58). The future establishment of this technique may considerably change therapeutic approaches to dental syndromes and diseases. However, there are some challenges before regeneration ie)Tooth embryogenesis takes longer duration and is highly-priced. There are more ethical and legal concerns like the morphology of the tooth might differ. We also need to further develop efficient protocols to induce stem cells to form cell types in vitro that are relevant to the tissues and organs targeted for regeneration. To succeed in these challenges, further basic studies to elucidate the regulatory mechanisms of stem cells and tooth development are needed.

ACKNOWLEDGMENT:

The authors are grateful to Saveetha Dental College for providing a platform to express our knowledge.

CONFLICT OF INTEREST:

No potential conflict of interest relevant to this article was reported.

REFERENCES:

1.Nanci A. Ten Cate's Oral Histology - E-Book: Development, Structure, and Function [Internet].ElsevierHealthSciences;2017.352p.Availablefrom:https://play.google.com/store/books/details?id=XR0xDwAAQBAJSciences;Availablefrom:

2. Angelova AV, Zaugg LK, Neves V, Liu Y, Sharpe PT, Others. Tooth Repair and Regeneration. Current oral health reports [Internet]. 2018;5(4):295–303. Available from: https://europepmc.org/articles/pmc6244610

3. Sarver DM, Proffit WR, Fields HW. Contemporary orthodontics [Internet]. Elsevier; 2007. Available from: http://125.234.102.146:8080/dspace/handle/DNULIB_52011/6101

4. Siddique R, Sureshbabu NM, Somasundaram J, Jacob B, Selvam D. Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi. J Conserv Dent [Internet]. 2019 Jan;22(1):40–7. Available from: http://dx.doi.org/10.4103/JCD.JCD_284_18

5. Rajakeerthi R, Ms N. Natural Product as the Storage medium for an avulsed tooth – A Systematic Review. 2019 Jun 11 [cited 2020 Jun 6];22(2):249–56. Available from: https://www.researchgate.net/publication/333706221_Natural_Product_as_the_Storage_medium_for_an_avulse d_tooth_-_A_Systematic_Review

6. Ramamoorthi S, Nivedhitha MS, Divyanand MJ. Comparative evaluation of postoperative pain after using endodontic needle and EndoActivator during root canal irrigation: A randomised controlled trial. Aust Endod J [Internet]. 2015 Aug;41(2):78–87. Available from: http://dx.doi.org/10.1111/aej.12076

7. Ramanathan S, Solete P. Cone-beam Computed Tomography Evaluation of Root Canal Preparation using Various Rotary Instruments: An in vitro Study. J Contemp Dent Pract [Internet]. 2015 Nov 1;16(11):869–72. Available from: http://dx.doi.org/10.5005/jp-journals-10024-1773

8. Nasim I, Hussainy S, Thomas T, Ranjan M. Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up [Internet]. Vol. 21, Journal of Conservative Dentistry. 2018. p. 510. Available from: http://dx.doi.org/10.4103/jcd.jcd_51_18

9. Kumar D, Delphine Priscilla Antony S. Calcified Canal and Negotiation-A Review. Research Journal of Pharmacy and Technology [Internet]. 2018 Aug 31 [cited 2020 Jun 6];11(8):3727–30. Available from: https://rjptonline.org/AbstractView.aspx?PID=2018-11-8-88

10. Pokorny PH, Wiens JP, Litvak H. Occlusion for fixed prosthodontics: a historical perspective of the gnathological influence. J Prosthet Dent [Internet]. 2008 Apr;99(4):299–313. Available from: http://dx.doi.org/10.1016/S0022-3913(08)60066-9

11. Jose J, P. A, Subbaiyan H. Different Treatment Modalities followed by Dental Practitioners for Ellis Class 2 Fracture – A Questionnaire-based Survey [Internet]. Vol. 14, The Open Dentistry Journal. 2020. p. 59–65. Available from: http://dx.doi.org/10.2174/1874210602014010059

12. Balic A. Biology Explaining Tooth Repair and Regeneration: A Mini-Review. Gerontology [Internet]. 2018 Mar 13;64(4):382–8. Available from: http://dx.doi.org/10.1159/000486592

13. Zhai Q, Dong Z, Wang W, Li B, Jin Y. Dental stem cell and dental tissue regeneration. Front Med [Internet]. 2019 Apr;13(2):152–9. Available from: http://dx.doi.org/10.1007/s11684-018-0628-x

14.Morsczeck C, Reichert TE. Dental stem cells in tooth regeneration and repair in the future. Expert Opin
BiolTher[Internet].2018Feb;18(2):187–96.Availablefrom:http://dx.doi.org/10.1080/14712598.2018.1402004

15. Oshima M, Mizuno M, Imamura A, Ogawa M, Yasukawa M, Yamazaki H, et al. Functional tooth regeneration using a bioengineered tooth unit as a mature organ replacement regenerative therapy. PLoS One [Internet]. 2011 Jul 12;6(7):e21531. Available from: http://dx.doi.org/10.1371/journal.pone.0021531

16. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study. Pesqui Bras Odontopediatria Clin Integr [Internet]. 2019 [cited 2020 Jun 6];19. Available from: https://www.scielo.br/scielo.php?pid=S1983-46322019000100364&script=sci_arttext

17. Nandakumar M, Nasim I. Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis. J Conserv Dent [Internet]. 2018 Sep;21(5):516–20. Available from: http://dx.doi.org/10.4103/JCD.JCD_110_18

18. Ravinthar K, Jayalakshmi. Recent Advancements in Laminates and Veneers in Dentistry. Research Journal of Pharmacy and Technology [Internet]. 2018 Feb 28 [cited 2020 Jun 6];11(2):785–7. Available from: https://rjptonline.org/AbstractView.aspx?PID=2018-11-2-70

19. Noor SSSE, S Syed Shihaab, Pradeep. Chlorhexidine: Its properties and effects [Internet]. Vol. 9, Research Journal of Pharmacy and Technology. 2016. p. 1755. Available from: http://dx.doi.org/10.5958/0974-360x.2016.00353.x

20. Teja KV, Ramesh S, Priya V. Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study. J Conserv Dent [Internet]. 2018 Nov;21(6):592–6. Available from: http://dx.doi.org/10.4103/JCD.JCD_154_18

21. Teja KV, Ramesh S. Shape optimal and clean more. Saudi Endodontic Journal [Internet]. 2019 Sep 1 [cited 2020 Jun 6];9(3):235. Available from: http://www.saudiendodj.com/article.asp?issn=1658-5984;year=2019;volume=9;issue=3;spage=235;epage=236;aulast=Teja 22. Mao AS, Mooney DJ. Regenerative medicine: Current therapies and future directions. Proc Natl Acad Sci U S A [Internet]. 2015 Nov 24;112(47):14452–9. Available from: http://dx.doi.org/10.1073/pnas.1508520112

23. Yu T, Volponi AA, Babb R, An Z, Sharpe PT. Stem Cells in Tooth Development, Growth, Repair, and Regeneration. Curr Top Dev Biol [Internet]. 2015 Oct 1;115:187–212. Available from: http://dx.doi.org/10.1016/bs.ctdb.2015.07.010

24. Slack JMW. Origin of stem cells in organogenesis. Science [Internet]. 2008 Dec 5;322(5907):1498–501. Available from: http://dx.doi.org/10.1126/science.1162782

25. Scadden DT. The stem-cell niche as an entity of action. Nature [Internet]. 2006 Jun 29;441(7097):1075–9. Available from: http://dx.doi.org/10.1038/nature04957

26. Li L, Tang Q, Wang A, Chen Y. Regrowing a tooth: in vitro and in vivo approaches. Curr Opin Cell Biol [Internet]. 2019 Dec;61:126–31. Available from: http://dx.doi.org/10.1016/j.ceb.2019.08.002

27. Volponi AA, Pang Y, Sharpe PT. Stem cell-based biological tooth repair and regeneration. Trends Cell Biol [Internet]. 2010 Dec;20(12):715–22. Available from: http://dx.doi.org/10.1016/j.tcb.2010.09.012

28. van Laake LW, Passier R, Doevendans PA, Mummery CL. Human embryonic stem cell-derived cardiomyocytes and cardiac repair in rodents. Circ Res [Internet]. 2008 May 9;102(9):1008–10. Available from: http://dx.doi.org/10.1161/CIRCRESAHA.108.175505

29. Alt-Holland A, Sowalsky AG, Szwec-Levin Y, Shamis Y, Hatch H, Feig LA, et al. Suppression of E-cadherin function drives the early stages of Ras-induced squamous cell carcinoma through upregulation of FAK and Src. J Invest Dermatol [Internet]. 2011 Nov;131(11):2306–15. Available from: http://dx.doi.org/10.1038/jid.2011.188

30. Otsu K, Kumakami-Sakano M, Fujiwara N, Kikuchi K, Keller L, Lesot H, et al. Stem cell sources for tooth regeneration: current status and future prospects. Front Physiol [Internet]. 2014 Feb 4;5:36. Available from: http://dx.doi.org/10.3389/fphys.2014.00036

31. Gronthos S, Mankani M, Brahim J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. Proc Natl Acad Sci U S A [Internet]. 2000 Dec 5;97(25):13625–30. Available from: http://dx.doi.org/10.1073/pnas.240309797

32. Janani K, Palanivelu A, Sandhya R. Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality - An in vivo study. 2020 Jan 31 [cited 2020 Jun 6];23(1). Available from: https://www.researchgate.net/publication/338959944_Diagnostic_accuracy_of_dental_pulse_oximeter_with_cu stomized_sensor_holder_thermal_test_and_electric_pulp_test_for_the_evaluation_of_pulp_vitality_-___An_in_vivo_study

33. Jo Y-Y, Lee H-J, Kook S-Y, Choung H-W, Park J-Y, Chung J-H, et al. Isolation and characterization of postnatal stem cells from human dental tissues. Tissue Eng [Internet]. 2007 Apr;13(4):767–73. Available from: http://dx.doi.org/10.1089/ten.2006.0192

34. Yamada Y, Ueda M, Naiki T, Takahashi M, Hata K-I, Nagasaka T. Autogenous injectable bone for regeneration with mesenchymal stem cells and platelet-rich plasma: tissue-engineered bone regeneration. Tissue Eng [Internet]. 2004 May;10(5-6):955–64. Available from: http://dx.doi.org/10.1089/1076327041348284

35. Huang GT-J, Gronthos S, Shi S. Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. J Dent Res [Internet]. 2009 Sep;88(9):792–806. Available from: http://dx.doi.org/10.1177/0022034509340867

36. Timothy CN, Samyuktha PS, Brundha MP. Dental pulp Stem Cells in Regenerative Medicine - A

Literature Review. J Adv Pharm Technol Res [Internet]. 2019;12(8):4052. Available from: http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=8&article=088

37. Nakamura S, Yamada Y, Katagiri W, Sugito T, Ito K, Ueda M. Stem cell proliferation pathways comparison between human exfoliated deciduous teeth and dental pulp stem cells by gene expression profile from promising dental pulp. J Endod [Internet]. 2009 Nov;35(11):1536–42. Available from: http://dx.doi.org/10.1016/j.joen.2009.07.024

38. Website [Internet]. [cited 2020 Jun 5]. Available from: Cordeiro MM, Dong Z, Kaneko T, Zhang Z, Miyazawa M, Shi S, et al. Dental pulp tissue engineering with stem cells from exfoliated deciduous teeth. J Endod [Internet]. 2008 Aug;34(8):962–9. Available from: http://dx.doi.org/10.1016/j.joen.2008.04.009

39. Shi S, Seo B-M, Miura M. Multipotent postnatal stem cells from human periodontal ligament and uses thereof [Internet]. US Patent. 9210925, 2015 [cited 2020 Jun 5]. Available from: https://patentimages.storage.googleapis.com/f4/89/4b/bab33f5337045d/US9210925.pdf

40. Yildirim S. Isolation Methods of Dental Pulp Stem Cells. In: Yildirim S, editor. Dental Pulp Stem Cells [Internet]. New York, NY: Springer New York; 2013. p. 41–51. Available from: https://doi.org/10.1007/978-1-4614-5687-2_5

41. Sonoyama W, Liu Y, Fang D, Yamaza T, Seo B-M, Zhang C, et al. Mesenchymal stem cell-mediated functional tooth regeneration in swine. PLoS One [Internet]. 2006 Dec 20;1:e79. Available from: http://dx.doi.org/10.1371/journal.pone.0000079

42. Huang GT-J, -J. Huang GT, Sonoyama W, Liu Y, Liu H, Wang S, et al. The Hidden Treasure in Apical Papilla: The Potential Role in Pulp/Dentin Regeneration and BioRoot Engineering [Internet]. Vol. 34, Journal of Endodontics. 2008. p. 645–51. Available from: http://dx.doi.org/10.1016/j.joen.2008.03.001

43. Coura GS, Garcez RC, de Aguiar CBNM, Alvarez-Silva M, Magini RS, Trentin AG. Human periodontal ligament: a niche of neural crest stem cells [Internet]. Vol. 43, Journal of Periodontal Research. 2008. p. 531–6. Available from: http://dx.doi.org/10.1111/j.1600-0765.2007.01065.x

44. Website [Internet]. [cited 2020 Jun 5]. Available from: Morsczeck C, Götz W, Schierholz J, Zeilhofer F, Kühn U, Möhl C, et al. Isolation of precursor cells (PCs) from human dental follicle of wisdom teeth. Matrix Biol [Internet]. 2005 Apr;24(2):155–65. Available from: http://dx.doi.org/10.1016/j.matbio.2004.12.004

45. Yao S, Pan F, Prpic V, Wise GE. Differentiation of stem cells in the dental follicle. J Dent Res [Internet]. 2008 Aug;87(8):767–71. Available from: http://dx.doi.org/10.1177/154405910808700801

46. Rincon JC, Young WG, Bartold PM. The epithelial cell rests of Malassez--a role in periodontal regeneration? J Periodontal Res [Internet]. 2006 Aug;41(4):245–52. Available from: http://dx.doi.org/10.1111/j.1600-0765.2006.00880.x

47. Xiong J, Gronthos S, Bartold PM. Role of the epithelial cell rests of Malassez in the development, maintenance and regeneration of periodontal ligament tissues. Periodontol 2000 [Internet]. 2013 Oct;63(1):217–33. Available from: http://dx.doi.org/10.1111/prd.12023

48. Yildirim O, Li R, Hung J-H, Chen PB, Dong X, Ee L-S, et al. Mbd3/NURD complex regulates expression of 5-hydroxymethylcytosine marked genes in embryonic stem cells. Cell [Internet]. 2011 Dec 23;147(7):1498–510. Available from: http://dx.doi.org/10.1016/j.cell.2011.11.054

49. Oshima M, Tsuji T. Whole Tooth Regeneration as a Future Dental Treatment. Adv Exp Med Biol [Internet]. 2015;881:255–69. Available from: http://dx.doi.org/10.1007/978-3-319-22345-2_14

50. Oshima M, Tsuji T. Whole Tooth Regeneration Using a Bioengineered Tooth. New Trends in Tissue Engineering and Regenerative Medicine Nagoya: IntechOpen [Internet]. 2014;109–19. Available from: https://books.google.co.in/books?hl=en&lr=&id=QGmQDwAAQBAJ&oi=fnd&pg=PA109&dq=oshima+et+al+

2011+tissue+engineering&ots=-Q6_5zDqK0&sig=pXHYqWFK28AZ1J8YRnOnrZGbbbA

51. Liu F, Chu EY, Watt B, Zhang Y, Gallant NM, Andl T, et al. Wnt/β-catenin signaling directs multiple stages of tooth morphogenesis. Dev Biol [Internet]. 2008 Jan 1;313(1):210–24. Available from: http://www.sciencedirect.com/science/article/pii/S0012160607014601

52. Park CH, Rios HF, Jin Q, Bland ME, Flanagan CL, Hollister SJ, et al. Biomimetic hybrid scaffolds for engineering human tooth-ligament interfaces. Biomaterials [Internet]. 2010 Aug;31(23):5945–52. Available from: http://dx.doi.org/10.1016/j.biomaterials.2010.04.027

53. Kim RY, Bae SS, Feinberg SE. Soft Tissue Engineering. Oral Maxillofac Surg Clin North Am [Internet]. 2017 Feb;29(1):89–104. Available from: http://dx.doi.org/10.1016/j.coms.2016.08.007

54. Valmikinathan CM, Tian J, Wang J, Yu X. Novel nanofibrous spiral scaffolds for neural tissue engineering. J Neural Eng [Internet]. 2008;5(4):422. Available from: https://iopscience.iop.org/article/10.1088/1741-2560/5/4/007/meta

55. Ma PX. Scaffolds for tissue fabrication. Mater Today [Internet]. 2004 May 1;7(5):30–40. Available from: http://www.sciencedirect.com/science/article/pii/S1369702104002330

56. Liu J, Yan C. 3D printing of scaffolds for tissue engineering. In: 3D Printing [Internet]. IntechOpen; 2018. Available from: https://books.google.co.in/books?hl=en&lr=&id=c1mRDwAAQBAJ&oi=fnd&pg=PA137&dq=Jingyu+Liu+et+ al+2018+internal+porous+structure&ots=6xx-EVqzNa&sig=k206YIZpUW13f-FF4QjFHHKkDmA

57. Young CS, Terada S, Vacanti JP, Honda M, Bartlett JD, Yelick PC. Tissue engineering of complex tooth structures on biodegradable polymer scaffolds. J Dent Res [Internet]. 2002 Oct;81(10):695–700. Available from: http://dx.doi.org/10.1177/154405910208101008

58. Manohar MP, Sharma S. A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists. Indian J Dent Res [Internet]. 2018 Nov;29(6):716–20. Available from: http://dx.doi.org/10.4103/ijdr.IJDR_716_16

59. Farhat Yaasmeen Sadique Basha, Rajeshkumar S, Lakshmi T, Anti-inflammatory activity of Myristica fragrans extract . Int. J. Res. Pharm. Sci., 2019 ;10(4), 3118-3120 DOI: https://doi.org/10.26452/ijrps.v10i4.1607